Crash cart supply & equipment checklist



Equipment
 □ Airway (oral and nasal) all sizes □ McGill forceps, large and small □ King Airway set (3) eliminates the need for laryngoscope and endotracheal tubes □ Bag valve mask (adult and pediatric) □ Nasal cannula □ Non-rebreather oxygen face masks (3 sizes) □ IV start packs □ Normal saline solution (1000ml bags) □ IV tubing □ Angiocaths (various sizes) □ 10ml normal saline flush syringes (3) □ Gauze □ Alcohol preps □ Monitor with defibrillator (preferred) or AED □ Syringe nasal adaptor (nasal Narcan atomizer)
Drugs □ Aspirin 81mg tablets (4) □ Nitroglycerin spray or 0.4mg tablets (3) □ Dextrose 50% (Dextrose 25% if treating pediatrics) □ Narcan 1mg/ml (6) □ Epinephrine 1:10,000 pre-filled emergency syringe (3) □ Atropine Sulfate 1mg pre-filled emergency syringe (3) □ Amiodarone 150mg vial (4) □ Epi Pen (2) □ Epi Pen (2) □ Epi Pen Jr (2) □ Solumedrol 125mg vial □ Benadryl 50mg Vial (2) □ Adenosine 6mg (3) □ Lopressor 10mg (2) □ Lidocaine 100mg/5mL pre-filled emergency syringe (2) □ Cardiazem 20mg vial (2) □ Pronestyl (Procainamide) 1g in 10 ml 100mg/ml vial (1)

Version control: This document follows 2020 American Heart Association® guidelines for CPR and ECC. American Heart Association® guidelines are updated every five years. If you are reading this page after December 2025, please contact support@acls.net for an update. Version 2024.11.a

please note that this checklist is a guideline, not an exhaustive list.